

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Books

Search PubMed



for



Limits

Preview/Index

History

Clipboard

Details

Display

Abstract



Show

20



Sort by



Send to



All: 1

Review: 0



About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI (Cubby)

Related Resources

Order Documents

NLM Mobile

NLM Catalog

NLM Gateway

TOXNET

Consumer Health


Clinical Alerts

ClinicalTrials.gov

PubMed Central

1: Neuroscience. 1995 Jan;64(2):385-95.

Related Articles. Links

**Alteration in nicotine binding sites in Parkinson's disease, Lewy body dementia and Alzheimer's disease: possible index of early neuropathology.****Perry EK, Morris CM, Court JA, Cheng A, Fairbairn AF, McKeith IG, Irving D, Brown A, Perry RH.**

MRC Neurochemical Pathology Unit, Newcastle General Hospital, Newcastle upon Tyne, U.K.

High-affinity nicotine binding, considered to primarily reflect the presence of CNS alpha 4 beta 2 nicotinic receptor subunits, was examined autoradiographically in brain regions most severely affected by Alzheimer and Parkinson types of pathology. In the midbrain, the high density of binding associated with the pars compacta of the substantia nigra was extensively reduced (65-75%, particularly in the lateral portion) in both Lewy body dementia and Parkinson's disease. Since loss of dopaminergic neurons in Lewy body dementia was only moderate (40%), loss or down-regulation of the nicotinic receptor may precede degeneration of dopaminergic neurons in this region. In the dorsolateral tegmentum, where diffuse cholinergic perikarya are located, nicotine binding was highly significantly decreased in both Lewy body dementia and Parkinson's disease with almost no overlap between the normal and disease groups, indicative of a major pathological involvement in or around the pedunculopontine cholinergic neurons. In the hippocampus, binding was decreased around the granular layer in Lewy body dementia and Alzheimer's disease, although unchanged in the stratum lacunosum moleculare, where binding was relatively higher. Dense bands of receptor binding in the presubiculum and parahippocampal gyrus--areas of highest binding in human cortex--were diminished in Alzheimer's disease but not Lewy body dementia. In temporal neocortex there were reductions in Alzheimer's disease throughout the cortical layers but in Lewy body dementia only in lower layers, in which Lewy bodies are concentrated. Abnormalities of the nicotinic receptor in the diseases examined appear to be closely associated with primary histopathological changes: dopaminergic cell loss in Parkinson's disease and Lewy body dementia, amyloid plaques and tangles in subicular and entorhinal areas in Alzheimer's disease. Loss or down-regulation of the receptor may precede neurodegeneration.

PMID: 7700528 [PubMed - indexed for MEDLINE]